

## BC ROUTINE IMMUNIZATION SCHEDULE INFANTS & CHILDREN

A G E	2 months	4 months	6 months	12 months	18 months	4-6 years
<b>VACCINE</b>						
<b>DTaP-HB-IPV-Hib</b> (diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b) <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>	✓	✓	✓			
<b>DTaP-IPV-Hib</b> <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>					✓	
<b>DTaP-IPV</b> (diphtheria, tetanus, pertussis, polio) <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>						✓
<b>Pneumococcal conjugate ‡</b> <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>	✓	✓		✓		
<b>Meningococcal conjugate C</b> <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>	✓			✓		
<b>MMR</b> (measles, mumps, rubella) <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>				✓	✓	
<b>Varicella</b> (chickenpox) <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a>				✓		
<b>Influenza</b> <a href="#">1. Vaccine HealthFile</a> <a href="#">2. Product Monograph</a> <a href="#">Vaxigrip</a> / <a href="#">Fluviral</a>			✓ *			

‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. See the web link for more information or speak to your doctor or public health nurse.

\* Second dose needed 4 weeks after the first if receiving vaccine for first time.